

## QUARANTINE COVID-19

<b>Move to Quarantine</b> <ul style="list-style-type: none"> <li>• New Intakes</li> <li>• Contacts</li> </ul>	Quarantine is used to separate <b>asymptomatic</b> persons who have <b>Risk Factors or who are Contacts to a person with COVID-19</b> during the incubation period (up to 14 days). Escort inmate, in mask, to a designated single room with door <b>OR</b> cohort with other asymptomatic inmates in a housing area with door. Staff escorting asymptomatic inmates with direct contact will wear at a minimum surgical mask, eye protection, and gloves.
<b>Implement Transmission Based Precautions - Standard/Contact/ Eye Protection/ Droplet.</b>  <b>PPE</b>	<b><i>Standard precautions/Contact/Eye Protection/Droplet Isolation</i></b> 1) Hand hygiene (before and after wearing gloves) 2) PPE ( <i>gloves, gown, eye protection, surgical mask</i> ) for entry into room and direct contact. 3) If not entering room and $\geq 6$ feet away, utilize standard precautions – gloves (e.g. place food container in food slot while inmate(s) stand at back of room). Inmate should wear mask or facial coverings, if able. - <b>Prior to room entry:</b> Perform hand hygiene. Apply (don) gloves, gown,* surgical mask and eye protection before room entry or inmate contact. <i>See donning checklist.</i> (*gown if available supply) - <b>Doffing with Anteroom:</b> Have inmate(s) move to a social distance $\geq 6$ feet, if possible, and remove gloves and gown* and then exit room. Perform hand hygiene, remove (doff) eye protection, mask and repeat hand hygiene. <b>IF no anteroom is available, exit out of room to doff all PPE in a designated doffing area (tape off area for doffing) located immediately outside of room.</b> <i>See doffing checklist. Deposit used PPE in a trash receptacle designated for this purpose near the doffing area.</i>
<b>Signage</b>	A Respiratory QUARANTINE Sign is placed on the door
<b>Inmate Education</b>	Advise/educate inmate regarding reportable symptoms of COVID-19 illness and notify housing unit office if symptoms arise. Educate regarding social distancing and facial covering. Provide education sheet.
<b>Communication and Documentation</b>	1) Notify facility leadership, infection prevention and control (QIIPC)/health services, Incident Command, Chief Psychologist and Regional QIIPC Consultants. 2) Conduct temperature checks twice daily, this can be conducted by non-healthcare staff after training. Any positive symptoms or febrile temperatures, move to isolation and notify health services staff (if screening conducted by non-health care staff) See Medical Care Below 3) Place a medical hold in BEMR and Sentry for the duration of the quarantine. Code inmate as (b)(7)(E); (b)(7)(F)
<b>Staff Interaction</b>	Staff assessments not requiring direct contact will be conducted with social distancing $\geq 6$ feet away. Limit the number of staff interactions with inmate(s) and take measures to reduce rotation of staff interacting with quarantined inmate(s). Dedicate personnel if possible.
<b>Medical Equipment</b>	Medical equipment should be dedicated to area if possible.
<b>Medical Care</b>	Isolate inmate(s) if symptomatic (cough, SOB, HA, dizziness, fatigue, loss of taste or smell, sore throat, chest pain) and/or a temperature $\geq 100.4$ F. Positive symptoms require Clinical Encounter. Limit close or direct contact as much as possible. Provide necessary medical care as needed.
<b>Food Service</b>	Regular trays or use disposable dish wear. Wear gloves and maintain social distancing. Dispose of in regular trash.
<b>Laundry</b>	Wear gloves. Regular central laundry processes are acceptable. Do not shake dirty laundry. Disinfect dirty carts after use.
<b>Visits</b>	In-person visits will be suspended until the end of quarantine. Consult local leadership for exceptions.
<b>Telephone Calls</b>	Phone should be cleaned and disinfected after each use with a product from the EPA registered disinfectant List N.
<b>Trash</b>	For disposal of trash, wear gloves and <b>double bag</b> in clear waste bags; <i>Ensure it is not processed by recycling.</i>
<b>Cleaning/Disinfection</b>	The inmate(s) should be provided supplies to clean room. Use disinfectant from EPA list N.
<b>Discontinuation of Quarantine</b>	Duration of quarantine is 14 days. <b>If at all possible</b> , do not add individuals to an existing quarantine after the 14-day quarantine clock has started, if no other option exists and new inmates are added into a quarantine cohort, the original group may be released from quarantine on the original schedule if no inmates develop COVID-19 symptoms or are diagnosed with COVID-19. Asymptomatic inmates should undergo COVID-19 testing resulting with negative findings prior to being released from quarantine.
<b>Terminal Cleaning</b>	When quarantine ends, the inmate(s), if possible, should clean the area. If inmates in quarantine became symptomatic, wait 24 hours (if possible), and then the area should be cleaned again with an EPA List N registered disinfectant while wearing gloves, gown and any other PPE recommended by the disinfectant manufacturer (i.e., if splashes are anticipated, wear mask and eye protection).